

On the difference between admissible and “differentiable” Hamiltonians

Vladimir O. Soloviev

Institute for High Energy Physics

Protvino, 142284, Moscow Region, Russia

e-mail: vosoloviev@mx.ihep.su

October 30, 1996

Abstract

It is shown that the Regge-Teitelboim criterion for fixing the unique boundary contribution to the Hamiltonian compatible with free boundary conditions should be modified if the Poisson structure is noncanonical. The new criterion requires cancellation of boundary contributions to the Hamiltonian equations of motion. In the same time, boundary contributions to the variation of Hamiltonian are allowed. The Ashtekar formalism for gravity and hydrodynamics of the ideal fluid with a free surface in the Clebsch variables are treated as examples.

PACS: 11.10.Ef, 04.20.Fy, 03.40.Gc

Keywords: Hamiltonian formalism, canonical gravity, Ashtekar variables, hydrodynamics, Clebsch variables, Poisson brackets, free boundary

In this publication we consider the case of free boundary conditions in field theory, i.e., the situation when variations of the field variables and their spatial derivatives are not necessarily equal to zero on the boundary of the relevant space domain. Therefore it would be incorrect to treat Hamiltonians (or Lagrangians) differing in spatial divergences as representatives of the same equivalence class. We are to fix a unique admissible Hamiltonian (or Lagrangian) from that class according to some rule.

The importance of such problems for physics can be easily seen from the long history of discussion on the role of surface integrals in the Hamiltonian of General Relativity [1] – [5]. This discussion had been resolved in a publication by Regge and Teitelboim [6] where a criterion for the adequate choice of divergences in the Hamiltonian was proposed. It was called the requirement of “differentiability” of the Hamiltonian (see also [7]).

Here we intend to show that this criterion, which is applicable for the canonical Poisson structure, should be revised in a more general situation, especially if boundary contributions appear in the symplectic form and (or) in the Poisson brackets. In the same time the general mathematical background staying behind the concrete formulation given to it by Regge and Teitelboim is untouched by our improvement.

This general background is the “natural boundary conditions” criterion of the variational calculus [8], [9]. The general idea of the natural boundary conditions is that we can get from a variational principle not only the Euler-Lagrange equations but also some equations on the boundary. Both of them follow from the requirement that the functional under study should have a stationary point. It is possible to derive the natural boundary conditions when arbitrary field variations on the boundary contribute to the functional variation, and so, their coefficients must be equal to zero to make the functional stationary.

The difference between admissible and “differentiable” Hamiltonians, which is discussed in the paper, arises when noncanonical variables are used in the Hamiltonian formalism. This situation is not unfamiliar. For example, noncanonical Poisson brackets may originate as Dirac brackets after some reduction procedure. Below we will consider two examples. The first one is the Ashtekar formulation of General Relativity where the noncanonicity of Poisson brackets arises as a result of the change of variables [10]. The second example is the Hamiltonian formalism for surface waves in ideal fluid. Here just a position of the boundary becomes a dynamical variable. In both cases

we will see that the relation between the Lagrangian and the Hamiltonian is not so simple as in the canonical situation. As a result we can see that the correspondence between the action functional and the boundary conditions survives in the general case whereas the requirement of “differentiability” for the Hamiltonian should be replaced by another one. The new criterion requires a cancellation of boundary terms in the Hamiltonian vector field, or saying it in other words, in the Hamiltonian equations of motion. The Hamiltonian vector field is to be constructed according to the so-called formal variational calculus [11] and its extension on divergences proposed in previous publications of the author [12], [13], [14].

Throughout this paper we will use the characteristic function of a relevant compact space domain Ω constructed by means of the Heaviside θ -function and smooth function $P(x)$ with the following properties

$$P(x) = \begin{cases} > 0 & \text{if } x \in \Omega; \\ = 0 & \text{if } x \in \partial\Omega; \\ < 0 & \text{otherwise.} \end{cases} \quad (1)$$

It allows us to write any integral over Ω formally as an integral over infinite space $\mathbb{R}^3$ and freely integrate by parts. We will omit d^3x in these integrals.

As a first example let us consider the Ashtekar formalism for canonical gravity [15]. It is well known that this formulation can be constructed as a result of field transformations starting from the tetrad variables in time gauge where the unit normal to spacelike hypersurface is taken as one leg of the tetrad [16]. The initial pair of canonical variables (E_{ia}, π^{ia}) can first be changed for another one $(\tilde{E}^{ia}, K_{ia})$

$$\{\tilde{E}^{ia}(x), K_j^b(y)\} = \frac{1}{2}\delta_j^i \delta^{ab} \delta(x, y), \quad (2)$$

where

$$\tilde{E}^{ia} = E E^{ia}, \quad K_i^a = K_{ij} E^{ja} + E^{-1} E_{ib} J^{ab}, \quad (3)$$

and $E_i^a E_a^j = \delta_i^j$, $E_i^a E_b^i = \delta_b^a$, $E = \det |E_{ia}|$, K_{ij} is the second fundamental form of the spacelike hypersurface, and the generator of triad (the three remaining vectors of the tetrad) rotations is

$$J^{ab} = \frac{1}{2}(K_i^a \tilde{E}^{bi} - K_i^b \tilde{E}^{ai}). \quad (4)$$

Then the second transformation is used which introduce the Ashtekar connection variable A_i^a instead of K_{ia}

$$A_i^a = iK_i^a + \Gamma_i^a, \quad \Gamma_i^a = \frac{1}{2}\epsilon^{abc}\tilde{E}_{jc}\tilde{E}^{jb}_{|i}, \quad (5)$$

where the vertical line denotes the standard Riemannian covariant derivative and

$$\{\tilde{E}^{ia}(x), A_j^b(y)\} = \frac{i}{2}\delta_j^i\delta^{ab}\delta(x, y). \quad (6)$$

This transformation leads to the new form of the internal rotation generator which we take as a primer of the admissible Hamiltonian

$$H(\hat{\lambda}^a) = 2 \int \theta(P) \hat{\lambda}^a \mathcal{D}_i \tilde{E}^{ia} \equiv 2i \int \theta(P) \hat{\lambda}^a \epsilon^{abc} J^{bc}. \quad (7)$$

Here the covariant derivative $\mathcal{D}_i$ is defined by the following relation

$$\mathcal{D}_i \xi^{ka} = \xi^{ka}_{|i} + \epsilon^{abc} A_i^b \xi^{kc}. \quad (8)$$

In contrast to its previous form given according to the formula (4), in the Ashtekar variables the generator density depends on the spatial derivatives of $\tilde{E}^{ia}$, and so its variation

$$\delta H = \int \left(\frac{\delta H}{\delta \tilde{E}^{ia}} \delta \tilde{E}^{ia} + \frac{\delta H}{\delta A_i^a} \delta A_i^a \right), \quad (9)$$

contains a boundary contribution

$$\frac{\delta H}{\delta \tilde{E}^{ia}} = -\theta_{,i} 2\hat{\lambda}^a - \theta 2\mathcal{D}_i \hat{\lambda}^a, \quad (10)$$

$$\frac{\delta H}{\delta A_i^a} = -\theta 2\epsilon^{abc} \hat{\lambda}^b \tilde{E}^{ic}. \quad (11)$$

If we deal with a free boundary condition case then this functional is not “differentiable” in the Regge-Teitelboim terminology. Now we will show that it is nevertheless admissible as it gives regular Hamiltonian equations.

The explanation follows from the fact that the Ashtekar variables are canonical only up to the boundary term [10]

$$\{A_i^a(x), A_j^b(y)\} = \theta_{,k} C^k_{[(ia)(jb)]} \delta(x, y), \quad (12)$$

where the square brackets denote antisymmetrization $i \leftrightarrow j, a \leftrightarrow b$ and

$$C_{(ia)(jb)}^k = \frac{i}{2E} (\epsilon^{acb} \delta_j^k E_{ic} - \epsilon^{acd} E_{ib} E_{jc} E^{kd}). \quad (13)$$

Due to this noncanonicity the Hamiltonian equations acquire a form

$$\dot{\tilde{E}}^{ia}(x) = \int \{ \tilde{E}^{ia}(x), A_j^b(y) \} \frac{\delta H}{\delta A_j^b(y)}, \quad (14)$$

$$\dot{A}_i^a(x) = \int \{ A_i^a(x), \tilde{E}^{jb}(y) \} \frac{\delta H}{\delta \tilde{E}^{jb}(y)} + \int \{ A_i^a(x), A_j^b(y) \} \frac{\delta H}{\delta A_j^b(y)}, \quad (15)$$

or, in the explicit form obtained by using the prescription from papers [12]-[14], (here it is simply $\theta(P) \cdot \theta'(P) = \theta'(P)$), they are

$$\dot{\tilde{E}}^{ia} = i\epsilon^{abc} \hat{\lambda}^c \tilde{E}^{ib}, \quad (16)$$

$$\dot{A}_i^a = i\mathcal{D}_i \hat{\lambda}^a, \quad (17)$$

where $\theta(P)$ -factors are omitted.

We can see that the singular on the boundary terms in the second equation have been mutually canceled despite their presence in the full variational derivative (10). It means that our Hamiltonian is admissible under arbitrary boundary conditions.

It is not surprising then that the Poisson algebra of these generators is closed irrespectable to boundary conditions

$$\{H(\hat{\xi}^a), H(\hat{\eta}^b)\} = H(i\epsilon^{cab} \hat{\xi}^a \hat{\eta}^b). \quad (18)$$

In the Lagrangian treatment it is evident that due to the absence of spatial derivatives in the action

$$S = 2 \int_{t_1}^{t_2} dt \int \theta(P) \left(K_i^a \dot{\tilde{E}}^{ia} - i\hat{\lambda}^a \epsilon^{abc} J^{bc} \right), \quad (19)$$

any boundary conditions are natural. The Ashtekar transformation gives it a new appearance

$$S = -2i \int_{t_1}^{t_2} dt \int \theta(P) \left((A_i^a - \Gamma_i^a) \dot{\tilde{E}}^{ia} - i\hat{\lambda}^a \mathcal{D}_i \tilde{E}^{ia} \right), \quad (20)$$

where this property is no more evident. But let us derive a variation of the action

$$\begin{aligned} \delta S = & -2i \int_{t_1}^{t_2} dt \int \theta(P) \left(\delta \tilde{E}^{ia} \left[-\dot{A}_i^a + i\mathcal{D}_i \tilde{E}^{ia} \right] + \delta A_i^a \left[\dot{\tilde{E}}^{ia} + i\varepsilon^{abc} \lambda^b \tilde{E}^{ic} \right] + \right. \\ & \left. + \frac{\partial}{\partial t} \left((A_i^a - \Gamma_i^a) \delta \tilde{E}^{ia} \right) + \left((\dot{\Gamma}_i^a \delta \tilde{E}^{ia} - \dot{\tilde{E}}^{ia} \delta \Gamma_i^a) - \partial_i (i\hat{\lambda}^a \delta \tilde{E}^{ia}) \right) \right). \quad (21) \end{aligned}$$

Then the total time derivative term gives zero contribution as field variations on the time interval boundary are zero, the terms in square brackets give the equations of motion and the rest terms give no contribution due to relation

$$(\dot{\Gamma}_i^a \delta \tilde{E}^{ia} - \dot{\tilde{E}}^{ia} \delta \Gamma_i^a) - \partial_i (i\hat{\lambda}^a \delta \tilde{E}^{ia}) = 0 \text{ mod } \left(\dot{\tilde{E}}^{ia} + i\varepsilon^{abc} \lambda^b \tilde{E}^{ic} = 0 \right), \quad (22)$$

valid for arbitrary functions $\delta \tilde{E}^{ia}$, that can be verified by the straightforward calculation.

As the second example we consider the Hamiltonian description of ideal fluid with a free surface. In Eulerian variables the action can be written [17] by means of Clebsch potentials

$$\mathbf{v} = \nabla \phi + \frac{\eta}{\rho} \nabla s + \frac{\beta}{\rho} \nabla \alpha, \quad (23)$$

in the following way

$$S = \int_{t_1}^{t_2} dt \int \theta(P) \left[\rho \left(\frac{\mathbf{v}^2}{2} - \Phi(\mathbf{x}) - \varepsilon(\rho, s) \right) - \rho \frac{D\phi}{Dt} - \eta \frac{Ds}{Dt} - \beta \frac{D\alpha}{Dt} - \tau K \right]. \quad (24)$$

where ρ is the fluid mass density, s is the specific entropy, $\varepsilon = \varepsilon(\rho, s)$ is the specific internal energy density, τ is the surface tension coefficient,

$$\frac{D}{Dt} = \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla, \quad K = -\nabla \cdot \left(\frac{\nabla P}{|\nabla P|} \right). \quad (25)$$

On the boundary the last formula gives the external curvature of it whereas ∇P is proportional to the normal, the proposal to use it is due to Abarbanel et al [18].

The corresponding symplectic form

$$- \int [\theta(P)(\delta\rho \wedge \delta\phi + \delta\eta \wedge \delta s + \delta\beta \wedge \delta\alpha) + \theta'(P)\delta P \wedge (\rho\delta\phi + \eta\delta s + \beta\delta\alpha)] . \quad (26)$$

is degenerate. To go to the Hamiltonian formalism we can use the Dirac procedure [19] or the Faddeev-Jackiw approach [20], but really both of them lead to the same result as a simple trick. If we introduce a canonical variable π , conjugate to P

$$S \rightarrow S + \int_{t_1}^{t_2} dt \int (\pi \dot{P} - \lambda\pi), \quad (27)$$

and so add to the symplectic form the standard term

$$\int \delta\pi \wedge \delta P, \quad (28)$$

then the modified symplectic form becomes nondegenerate and can be inverted for getting the Poisson bivector

$$\begin{aligned} \Psi = & \int \left[\theta(P) \left(\frac{\delta}{\delta\rho} \wedge \frac{\delta}{\delta\phi} + \frac{\delta}{\delta\eta} \wedge \frac{\delta}{\delta s} + \frac{\delta}{\delta\beta} \wedge \frac{\delta}{\delta\alpha} \right) + \right. \\ & \left. + \frac{\delta}{\delta P} \wedge \frac{\delta}{\delta\pi} + \theta'(P) \frac{\delta}{\delta\pi} \wedge \left(\rho \frac{\delta}{\delta\rho} + \eta \frac{\delta}{\delta\eta} + \beta \frac{\delta}{\delta\beta} \right) \right], \end{aligned} \quad (29)$$

which explicitly contains the boundary δ -function.

The Hamiltonian has the following form

$$H = \int \left[\theta(P) \left(\frac{\rho \mathbf{v}^2}{2} + \rho\Phi + \rho\varepsilon(\rho, s) + \tau K \right) + \lambda\pi \right], \quad (30)$$

and its variation also have singular boundary contributions, for example,

$$\frac{\delta H}{\delta P} = \theta'(P) \left(\frac{\rho \mathbf{v}^2}{2} + \rho\Phi + \rho\varepsilon + \tau K \right), \quad (31)$$

$$\frac{\delta H}{\delta\phi} = -\theta'(P)\rho\mathbf{v} \cdot \nabla P - \theta(P)\nabla(\rho\mathbf{v}), \quad (32)$$

and so on.

Let us estimate the Hamiltonian vector field according to the standard formula where the interior product should be understood according to the definition given in Refs. [13], [14]

$$\begin{aligned}
- \mathbf{dH} \lrcorner \Psi &= [-\theta'(P)\rho(\lambda + \mathbf{v} \cdot \nabla P) - \theta(P)\nabla(\rho\mathbf{v})] \frac{\delta}{\delta\rho} + \\
&+ [-\theta'(P)\eta(\lambda + \mathbf{v} \cdot \nabla P) - \theta(P)(\nabla(\eta\mathbf{v}) - \rho T)] \frac{\delta}{\delta\eta} + \\
&+ [-\theta'(P)\beta(\lambda + \mathbf{v} \cdot \nabla P) - \theta(P)\nabla(\beta\mathbf{v})] \frac{\delta}{\delta\beta} + \\
&+ \theta'(P)(p - \tau K) \frac{\delta}{\delta\pi} + \\
&+ \theta(P) \left[\frac{v^2}{2} - \mathbf{v} \cdot \nabla\phi - \Phi - \varepsilon - \frac{p}{\rho} \right] \frac{\delta}{\delta\phi} - \theta(P) [\mathbf{v} \cdot \nabla s] \frac{\delta}{\delta s} - \\
&- \theta(P)\mathbf{v} \cdot \nabla\alpha \frac{\delta}{\delta\alpha} + \theta(P)\lambda \frac{\delta}{\delta P}, \tag{33}
\end{aligned}$$

here $p = \rho^2 \frac{\partial\varepsilon}{\partial\rho}$ is the pressure and $T = \frac{\partial\varepsilon}{\partial s}$ is the temperature.

Then the requirement that this Hamiltonian vector field must not contain boundary terms is equivalent to the standard boundary conditions for the problem [21]

$$\theta'(P)(\dot{P} + \mathbf{v} \cdot \nabla P) = 0, \tag{34}$$

$$\theta'(P)(p - \tau K) = 0, \tag{35}$$

if we take into account equation of motion

$$\dot{P} = \lambda. \tag{36}$$

It is useful to compare this approach with the analysis made in other variables [18].

If we try to use the criterion by Regge and Teitelboim here we will get the wrong boundary conditions, for example, one of them will be

$$\mathbf{v} \cdot \nabla P = 0, \tag{37}$$

that is, the requirement for the boundary to be fixed.

Lagrangian approach to the problem consists in studying the variation of the action (24). Apart from terms giving equations of motion this variation also contains boundary terms

$$\begin{aligned} \delta' S = & \int_{t_1}^{t_2} dt \int \left(-\frac{\partial}{\partial t} (\theta \rho \delta \phi + \theta \eta \delta s + \theta \beta \delta \alpha) + \right. \\ & \left. + \theta' (\dot{P} + \mathbf{v} \cdot \nabla P) (\rho \delta \phi + \eta \delta s + \beta \delta \alpha) + \theta' (p - \tau K) \delta P \right). \end{aligned} \quad (38)$$

The total time derivative gives the symplectic form (26) and does not contribute to the variational principle because the field variations are zero on the time boundary. The other terms contribute at the spatial boundary where the field variations are arbitrary and just give us the natural boundary conditions which are the same as (34), (35).

We have shown that in general situation when the Hamiltonian variables are not necessarily canonical and their Poisson brackets may contain boundary terms the Regge-Teitelboim criterion of “differentiability” of the Hamiltonian must be replaced by a new one. A Hamiltonian may be considered as admissible if the Hamiltonian vector field constructed according to the extended definitions given in [13], [14], does not contain any boundary contribution. This completes the search for a new criterion started in [10]. In general form the idea can be also traced to a publication by Mason [22]. More detailed treatment will be given elsewhere.

We hope that the Hamiltonian approach to field theory with free boundary conditions will be useful in dealing with different physical problems, especially those where the Lagrangian approach meets with difficulties.

Acknowledgements

This work has been completed during a visit of the author to the International Centre for Theoretical Physics in Trieste. The author is most grateful to Professor S. Randjbar-Daemi for the invitation and kind hospitality, partial support from ICTP is gratefully acknowledged. It is a pleasure to thank A. Zheltukhin for discussion.

References

- [1] R. Arnowitt, S. Deser and Ch.W. Misner, in: Gravitation, an Introduction to Current Research, ed. L. Witten (New York, 1963).
- [2] P.A.M. Dirac, Phys. Rev. Lett. 2 (1959) 368; Phys. Rev. 114 (1959) 924.
- [3] P.W. Higgs, Phys. Rev. Lett. 3 (1959) 66.
- [4] J. Schwinger, Phys. Rev. 130 (1963) 1253.
- [5] B.S. DeWitt, Phys. Rev. 160 (1967) 1113.
- [6] T. Regge and C. Teitelboim, Ann. Phys. 88 (1974) 286.
- [7] L.A. Takhtajan, L.D. Faddeev, Hamiltonian Method in Soliton Theory, (Springer-Verlag, N.Y., 1987) pp.18-19.
- [8] R. Courant and D. Hilbert, Methods of Mathematical Physics, Vol.1, (Wiley, N.Y., 1989) pp.208-211.
- [9] C. Lanczos, The Variational Principles of Mechanics, (Univ. Toronto Press, 1964) pp.68-73.
- [10] V.O. Soloviev, Phys. Lett. B 292 (1992) 30.
- [11] I.M. Gel'fand and L.A. Dickey, Uspekhi Matematicheskikh Nauk 30 (1975) 67.
- [12] V.O. Soloviev, J. Math. Phys. 34 (1993) 5747.
- [13] V.O. Soloviev, Boundary values as Hamiltonian variables II. Graded structures, Preprint IHEP 94-145, Protvino, 1994, q-alg/9501017 (submitted to J. Math. Phys.).
- [14] V.O. Soloviev, Nucl. Phys. BP49 (1996) 35.

- [15] A. Ashtekar, Phys. Rev. Lett. 57 (1986) 2244; Phys. Rev. D36 (1987) 1587; A. Ashtekar, P. Mazur, C.G. Torre, Phys. Rev. D36 (1987) 2955.
- [16] M. Henneaux, J.E. Nelson, C. Schomblond, Phys. Rev. D39 (1989) 434.
- [17] R.L. Seliger and G.B. Whitham, Proc. Roy. Soc. A305 (1968) 1.
- [18] H.D.I. Abarbanel, R. Brown and Y.M. Yang, Phys. Fluids 31 (1988) 2802.
- [19] P. Dirac, Lectures on Quantum Mechanics, (Yeshiva Univ., N.Y., 1964).
- [20] L. Faddeev and R. Jackiw, Phys. Rev. Lett. 60 (1988) 1692.
- [21] H. Lamb, Hydrodynamics, (Cambridge Univ. Press, Cambridge, 1932) p.456.
- [22] L. Mason, Class. Quant. Grav. 6 (1989) L7.